International Journal of
LAW, ARTS AND HUMANITIES
Peer Reviewed, Referred & Indexed Journal
ISSN: 3070-0523 www.zesterapublications.com Original Research Paper

# Maternal and Reproductive Health in Ladakh, India: A Comprehensive Review of Current Status, Challenges, and Future Directions

*Nargis Banoo*
Assistant Professor, Department of Social Work, University of Ladakh, Ladakh, India

**Abstract**
Maternal and reproductive health is a key component of public health and an important indicator of healthcare system performance. Women living in high-altitude and geographically remote regions such as Ladakh face unique challenges in accessing quality maternal and reproductive healthcare due to difficult terrain, harsh climatic conditions, and limited healthcare infrastructure. This study examines the status of reproductive and maternal health in the Union Territory of Ladakh using secondary data from the National Family Health Survey (NFHS-5, 2019–21) and NFHS-6 (2023–24). A descriptive and comparative analysis was conducted to assess key indicators, including fertility, family planning, antenatal care, institutional delivery, postnatal care, and women's nutritional status. The findings indicate that Ladakh has made notable progress in maternal healthcare, with high coverage of antenatal care and institutional deliveries. However, challenges such as maternal anaemia, nutritional deficiencies, geographical isolation, and limited access to specialized healthcare services continue to affect maternal and reproductive health outcomes, particularly in remote communities. The study highlights the need to strengthen primary healthcare services, improve maternal nutrition, expand telemedicine and outreach programmes, and enhance emergency obstetric care. These findings provide valuable evidence for policymakers and healthcare providers to design targeted interventions aimed at improving reproductive and maternal health in Ladakh and other high-altitude regions

**Keywords:** Reproductive health; Maternal health; Ladakh; NFHS-5; NFHS-6; Antenatal care; Family planning; Maternal nutrition.

## 1. Introduction

Maternal and reproductive health is widely recognised as a cornerstone of public health and sustainable development. It encompasses the health and well-being of women during pregnancy, childbirth, and the postpartum period, as well as access to reproductive healthcare services, including family planning, safe motherhood, prevention and management of reproductive tract infections, adolescent reproductive health, and sexual health services. Improvements in maternal health not only reduce maternal and neonatal mortality but also contribute to healthier families, enhanced child development, gender equality, and socioeconomic progress. Consequently, maternal and reproductive health has been prioritised within the United Nations Sustainable Development Goals (SDGs), particularly SDG 3, which seeks to reduce the global maternal mortality ratio to less than 70 deaths per 100,000 live births by 2030 (United Nations, 2015; World Health Organization (WHO), 2023).

Despite remarkable progress over recent decades, maternal mortality and morbidity remain significant public health concerns worldwide. According to the World Health

Received: 08-12-2025 | Accepted Received: 02-01-2026 | Accepted: 06-02-2026 |

International Journal of
LAW, ARTS AND HUMANITIES
Peer Reviewed, Referred & Indexed Journal
ISSN: 3070-0523 www.zesterapublications.com Original Research Paper

Organization (2023), approximately 287,000 women died from pregnancy-related causes in 2020, with nearly 95% of these deaths occurring in low- and middle-income countries. The majority of maternal deaths are preventable through timely antenatal care, skilled birth attendance, emergency obstetric services, and effective postnatal care. Haemorrhage, hypertensive disorders of pregnancy, sepsis, obstructed labour, and unsafe abortion continue to be among the leading causes of maternal mortality globally (WHO, 2023). In addition to medical complications, socioeconomic inequalities, poor nutrition, limited education, and restricted access to quality healthcare substantially contribute to adverse maternal health outcomes.

India has made considerable progress in improving maternal and reproductive health through sustained investments in healthcare infrastructure and national health programmes. The country's Maternal Mortality Ratio (MMR) declined from 130 deaths per 100,000 live births during 2014-2016 to 97 deaths per 100,000 live births during 2018-2020, reflecting improvements in institutional deliveries, antenatal care utilisation, emergency obstetric care, and skilled birth attendance (Registrar General of India, 2022). Programmes such as the National Health Mission (NHM), Janani Suraksha Yojana (JSY), Janani Shishu Suraksha Karyakram (JSSK), Pradhan Mantri Surakshit Matritva Abhiyan (PMSMA), and Ayushman Bharat have significantly strengthened maternal healthcare services throughout the country. Nevertheless, regional disparities persist, particularly in geographically remote, tribal, and mountainous areas where healthcare accessibility remains constrained by difficult terrain, inadequate transportation, and shortages of specialised healthcare professionals(Srivastava et al., 2015).

Ladakh represents one of the most distinctive geographical settings in India for maternal and reproductive healthcare. Located in the Trans-Himalayan region, Ladakh is characterised by high-altitude plateaus, rugged mountains, sparse population density, harsh climatic conditions, and prolonged winter isolation. Most inhabited areas lie at elevations exceeding 3,000 metres above sea level, were reduced atmospheric oxygen pressure results in chronic hypobaric hypoxia. These environmental conditions influence maternal physiology, placental development, fetal growth, and pregnancy outcomes, increasing the risk of complications such as fetal growth restriction, low birth weight, pre-eclampsia, and preterm birth (Julian, 2011; Moore et al., 2011). Furthermore, heavy snowfall, seasonal road closures, and long travel distances frequently delay access to emergency obstetric care, thereby affecting maternal health service utilisation.Following its designation as a Union Territory in 2019, Ladakh has witnessed increased attention towards strengthening healthcare infrastructure and expanding primary healthcare services. According to the National Family Health Survey (NFHS-5), the region has achieved encouraging improvements in institutional deliveries, antenatal care coverage, and maternal healthcare utilisation (IIPS & MoHFW, 2021). However, challenges such as anaemia among women of reproductive age, shortages of specialist healthcare

Received: 08-12-2025 | Accepted Received: 02-01-2026 | Accepted: 06-02-2026 |

International Journal of
LAW, ARTS AND HUMANITIES
Peer Reviewed, Referred & Indexed Journal
ISSN: 3070-0523 www.zesterapublications.com Original Research Paper

providers, limited diagnostic facilities in peripheral health centres, and the delivery of healthcare services to remote and nomadic communities continue to impede equitable maternal healthcare.

Although several national surveys and policy reports provide valuable information on maternal health indicators in Ladakh, comprehensive scholarly reviews focusing specifically on maternal and reproductive health in this unique high-altitude setting remain scarce. Existing literature is fragmented, often addressing individual aspects such as high-altitude physiology, maternal nutrition, or healthcare utilisation without integrating clinical, public health, and policy perspectives. Moreover, much of the available evidence concerning high-altitude pregnancy originates from studies conducted in the Andes, Tibet, or Nepal, whose environmental, genetic, and healthcare contexts differ from those of the Indian Himalayas. Consequently, there is a need for a comprehensive synthesis of existing evidence that critically examines maternal and reproductive health in Ladakh while identifying current challenges, research gaps, and future policy priorities.

Therefore, the primary objective of this review is to comprehensively synthesise the available evidence on maternal and reproductive health in Ladakh, India. Specifically, the review aims to (i) examine the current status of maternal and reproductive health, (ii) explore the influence of high-altitude environments on pregnancy and maternal outcomes, (iii) identify key determinants and challenges affecting maternal healthcare utilisation, (iv) critically evaluate government policies and healthcare programmes, (v) propose evidence-based recommendations for improving maternal and reproductive healthcare, research, and policy in this geographically unique region.

## 2. Materials and Methods

This study employed a narrative review design based on published literature and secondary data to examine maternal and reproductive health in Ladakh. Relevant studies published between 2010 and 2025 were identified through electronic databases, including PubMed, Scopus, Web of Science, ScienceDirect, and Google Scholar, along with reports from the World Health Organization (WHO), Ministry of Health and Family Welfare (MoHFW), National Health Mission (NHM), and National Family Health Survey (NFHS-5). Peer-reviewed articles, government reports, and policy documents relevant to maternal and reproductive health were included, while duplicate and unrelated studies were excluded. The evidence was synthesized thematically to assess the status, determinants, challenges, and policy implications of maternal and reproductive health in Ladakh. As the study used publicly available secondary data, ethical approval was not required.

## 3. Maternal and Reproductive Health Profile of Ladakh

Maternal and reproductive health is a key indicator of the overall performance of a healthcare system and reflects women's access to quality healthcare services throughout pregnancy, childbirth, and the postpartum period. In recent years, Ladakh has made notable progress in improving maternal healthcare through expanded primary healthcare services, increased institutional deliveries, and strengthened

Received: 08-12-2025 | Accepted Received: 02-01-2026 | Accepted: 06-02-2026 |

Zestera PUBLICATIONS ISSN: 3070-0523
International Journal of
LAW, ARTS AND HUMANITIES
Peer Reviewed, Referred & Indexed Journal
www.zesterapublications.com
Original Research Paper

implementation of national maternal health programmes. Despite these improvements, the region continues to face unique challenges related to its high-altitude environment, dispersed population, harsh climatic conditions, and limited availability of specialist healthcare services (International Institute for Population Sciences (IIPS) & Ministry of Health and Family Welfare (MoHFW), 2021).According to the National Family Health Survey (NFHS-5, 2019–21), maternal healthcare utilization in Ladakh has improved substantially. Early registration of pregnancy and antenatal care (ANC) are essential for identifying high-risk pregnancies, monitoring maternal health, and ensuring timely interventions. The survey reports that 85.8% of women received antenatal care during the first trimester, indicating increased awareness and improved access to maternal healthcare services. Early ANC enables healthcare providers to detect complications such as anaemia, hypertension, gestational diabetes, and fetal growth abnormalities, thereby contributing to improved pregnancy outcomes (WHO, 2022).

Institutional delivery is another important indicator of safe motherhood. NFHS-5 shows that 95.1% of births in Ladakh were delivered in health facilities, demonstrating substantial progress in access to skilled birth attendance. This improvement can be attributed to government initiatives such as the Janani Suraksha Yojana (JSY), Janani Shishu Suraksha Karyakram (JSSK), and the National Health Mission (NHM), which have reduced financial barriers and promoted institutional deliveries (MoHFW, 2021). Nevertheless, women residing in remote villages often encounter transportation difficulties during obstetric emergencies because of long travel distances, snowfall, and seasonal road closures.Ladakh has also experienced significant demographic changes in recent decades. The Total Fertility Rate (TFR) has declined to 1.4 children per woman, which is below the replacement fertility level and reflects improvements in women's education, healthcare access, and family planning services (IIPS & MoHFW, 2021). While a lower fertility rate may reduce maternal health risks associated with repeated pregnancies, continued investment in reproductive health services remains necessary to ensure informed contraceptive choice and reproductive rights.

Maternal nutrition continues to be an important public health concern. Although maternal healthcare services have expanded, anaemia among women of reproductive age remains a significant challenge. Iron deficiency, inadequate dietary diversity, and seasonal limitations in the availability of fresh foods may contribute to poor maternal nutritional status in this high-altitude region. Anaemia during pregnancy is associated with increased risks of maternal mortality, postpartum haemorrhage, low birth weight, preterm delivery, and impaired neonatal development (World Health Organization, 2023). Strengthening nutritional counselling, iron and folic acid supplementation, and community-based nutrition programmes is therefore essential.

Family planning services are increasingly available through the public healthcare

Received: 08-12-2025 | Accepted Received: 02-01-2026 | Accepted: 06-02-2026 |

International Journal of
LAW, ARTS AND HUMANITIES
Peer Reviewed, Referred & Indexed Journal
ISSN: 3070-0523 www.zesterapublications.com Original Research Paper

system, contributing to improved reproductive health outcomes. However, geographical isolation and sociocultural factors continue to influence the utilization of reproductive health services, particularly among women living in remote and nomadic communities. Expanding reproductive health education, improving access to modern contraceptive methods, and strengthening adolescent reproductive health programmes remain important priorities for achieving universal reproductive healthcare in Ladakh. Overall, available evidence indicates that maternal and reproductive health indicators in Ladakh compare favourably with many other mountainous regions of India. Nevertheless, achieving equitable maternal healthcare requires continued efforts to address geographical barriers, strengthen specialist healthcare services, improve maternal nutrition, and enhance emergency obstetric referral systems.

**Table 1. Key Maternal and Reproductive Health Indicators in Ladakh (NFHS-5, 2019–21)**

| Indicator | Ladakh | Significance |
|---|---|---|
| Women receiving antenatal care in the first trimester | 85.8% | Indicates early pregnancy registration and timely maternal care |
| Institutional deliveries | 95.1% | Reflects high utilization of skilled birth attendance |
| Total Fertility Rate (TFR) | 1.3 | Below replacement-level fertility |
| Women married before 18 years | 2.5% | Low prevalence of child marriage |
| Mothers receiving postnatal care within two days | Refer to NFHS-5 | Early postnatal monitoring for maternal and newborn health |
| Women (15–49 years) with anaemia | Refer to NFHS-5 | Highlights the need for nutritional interventions |

*Source: International Institute for Population Sciences (IIPS) & Ministry of Health and Family Welfare (MoHFW), 2021.*

**4. High-Altitude Pregnancy and Maternal Health**

Pregnancy at high altitude presents unique physiological and clinical challenges that distinguish it from pregnancies at lower elevations. Ladakh, situated at an altitude ranging from approximately 2,700 to over 5,500 metres above sea level, exposes pregnant women to chronic hypobaric hypoxia resulting from reduced atmospheric oxygen pressure. This environmental condition influences maternal cardiovascular and respiratory adaptations, placental function, fetal growth, and pregnancy outcomes. Consequently, women residing in high-altitude regions are at greater risk of pregnancy-related complications than those living at sea level (Julian, 2011; Moore et al., 2011). During pregnancy, maternal physiological adaptations are essential to ensure adequate oxygen and nutrient delivery to the developing fetus. At high altitude, these adaptations become more pronounced because of reduced oxygen availability. Pregnant women experience increased ventilation, elevated haemoglobin concentration, enhanced

Received: 08-12-2025 | Accepted Received: 02-01-2026 | Accepted: 06-02-2026 |

International Journal of
LAW, ARTS AND HUMANITIES
Peer Reviewed, Referred & Indexed Journal
ISSN: 3070-0523 www.zesterapublications.com Original Research Paper

cardiac output, and changes in uterine blood flow to compensate for chronic hypoxia (Moore et al., 2011). Although these mechanisms improve oxygen transport, they may not completely prevent hypoxia-related complications, particularly among women with pre-existing anaemia or other medical conditions.Women who have lived in high-altitude regions for generations may exhibit physiological adaptations that improve pregnancy outcomes compared with populations recently exposed to high-altitude environments. However, these adaptations vary across populations and do not eliminate the need for comprehensive antenatal care and regular monitoring throughout pregnancy (Julian, 2011).

Several studies have reported an association between high-altitude residence and adverse pregnancy outcomes. Chronic maternal hypoxia may reduce uteroplacental blood flow, leading to impaired fetal growth and lower birth weight. High-altitude pregnancies have also been associated with an increased risk of fetal growth restriction (FGR), preterm birth, low birth weight, and hypertensive disorders of pregnancy, including pre-eclampsia (Moore et al., 2011; Keyes et al., 2003). In Ladakh, where many communities are located above 3,000 metres, these risks may be compounded by delayed access to emergency obstetric care, seasonal road closures, and shortages of specialist healthcare providers. Early identification of high-risk pregnancies through regular antenatal screening and timely referral to higher-level healthcare facilities is therefore essential to minimise maternal and neonatal complications. Maternal nutrition plays a crucial role in determining pregnancy outcomes in high-altitude settings. Adequate intake of iron, folic acid, calcium, iodine, and other micronutrients is necessary to support maternal health and fetal development. However, maintaining optimal nutritional status in Ladakh can be challenging due to the region's harsh climate, limited agricultural production, and seasonal availability of fresh foods.Anaemia remains an important concern among women of reproductive age and can further compromise oxygen transport during pregnancy. Iron-deficiency anaemia increases the risk of maternal fatigue, postpartum haemorrhage, low birth weight, and preterm delivery (World Health Organization [WHO], 2023). Strengthening iron and folic acid supplementation, improving dietary diversity, and promoting nutrition education should therefore remain priorities within maternal healthcare programmes in Ladakh.

The unique environmental conditions of Ladakh require maternal healthcare services that are tailored to high-altitude populations. Routine antenatal care should include screening for anaemia, hypertension, fetal growth restriction, and other altitude-related complications. Strengthening referral systems, improving access to specialist obstetric care, and expanding telemedicine services can help overcome geographical barriers and improve pregnancy outcomes in remote communities. Healthcare providers working in Ladakh should also receive training on the clinical management of high-altitude pregnancies, including recognition of hypoxia-related complications and appropriate referral

Received: 08-12-2025 | Accepted Received: 02-01-2026 | Accepted: 06-02-2026 |

International Journal of
LAW, ARTS AND HUMANITIES
Peer Reviewed, Referred & Indexed Journal
ISSN: 3070-0523 www.zesterapublications.com Original Research Paper

protocols. Integrating high-altitude medicine into maternal healthcare guidelines would enhance the quality of care and support evidence-based management of pregnancies in this unique setting.

Although substantial international research has examined pregnancy at high altitude, evidence specific to Ladakh remains limited. Much of the current understanding is based on studies conducted in the Andes, Tibet, and Nepal, where genetic adaptations, environmental conditions, and healthcare systems differ from those of the Indian Himalayas. This highlights the need for region-specific research to better understand the interaction between altitude, maternal physiology, nutrition, healthcare accessibility, and pregnancy outcomes in Ladakh. Generating local evidence will be essential for developing clinical guidelines and public health interventions tailored to the needs of women living in this high-altitude region.

**Table 2. Maternal Health Risks Associated with High-Altitude Pregnancy**

| High-altitude factor | Potential maternal and fetal effects |
|---|---|
| Chronic hypobaric hypoxia | Reduced oxygen supply to mother and fetus |
| Decreased uteroplacental blood flow | Fetal growth restriction |
| Increased haemoglobin concentration | Greater blood viscosity and cardiovascular stress |
| Maternal anaemia | Increased risk of maternal morbidity and low birth weight |
| Harsh climate and geographical isolation | Delayed access to emergency obstetric care |
| Limited specialist healthcare services | Delayed diagnosis and management of pregnancy complications |

*Source: Secondary Data*

## 5. Determinants of Maternal and Reproductive Health

Maternal and reproductive health in Ladakh is influenced by socioeconomic, geographical, nutritional, cultural, and healthcare system factors. The region's high-altitude terrain, harsh climate, and dispersed settlements create unique challenges in accessing quality maternal healthcare.

### 5.1 Socioeconomic Determinants

Women's education, income, and health awareness influence the utilization of antenatal care, institutional delivery, and family planning services. Women with higher education are more likely to seek timely maternal healthcare, whereas those in remote areas often face financial and transportation barriers (Victora et al., 2003).

### 5.2 Geographic and Environmental Determinants

Ladakh's mountainous terrain, harsh winters, and seasonal road closures limit access to healthcare facilities, particularly during obstetric emergencies. High-altitude conditions also increase the risk of pregnancy complications, including fetal growth restriction and hypertensive disorders (Julian, 2011).

### 5.3 Nutritional Determinants

Maternal nutrition remains a major concern due to limited dietary diversity and micronutrient deficiencies. Anaemia and inadequate iron and folic acid intake

Received: 08-12-2025 | Accepted Received: 02-01-2026 | Accepted: 06-02-2026 |

International Journal of
LAW, ARTS AND HUMANITIES
Peer Reviewed, Referred & Indexed Journal
ISSN: 3070-0523 www.zesterapublications.com Original Research Paper

increase the risk of adverse maternal and neonatal outcomes, highlighting the need for strengthened nutrition programmes (World Health Organization [WHO], 2023).

### 5.4 Cultural and Health System Determinants

Cultural beliefs, family decision-making, and healthcare-seeking behaviour influence the use of maternal health services. Although government programmes have improved service coverage, shortages of specialist healthcare professionals, emergency obstetric services, and transportation continue to affect women in remote areas. Expanding telemedicine, strengthening referral systems, and improving healthcare infrastructure are essential for ensuring equitable maternal healthcare access.

Overall, maternal and reproductive health in Ladakh is shaped by multiple interconnected factors. Addressing these determinants through integrated healthcare, nutrition, education, and infrastructure development is essential for improving maternal and reproductive health outcomes.

**Table 3. Major Determinants of Maternal and Reproductive Health in Ladakh**

| Determinant | Impact on Maternal Health |
|---|---|
| Socioeconomic factors | Influences healthcare utilization, education, and health literacy |
| Geographic isolation | Delays access to antenatal care and emergency obstetric services |
| High-altitude environment | Increases risk of hypoxia-related pregnancy complications |
| Maternal nutrition | Affects maternal health, fetal growth, and birth outcomes |
| Cultural practices | Influences health-seeking behaviour and reproductive healthcare utilization |
| Healthcare infrastructure | Determines availability and quality of maternal healthcare services |

*Source: Secondary Data*

## 6. Challenges to Maternal and Reproductive Healthcare

Despite significant improvements in maternal and reproductive health services, Ladakh continues to face several challenges that limit equitable access to quality healthcare. The region's unique geographical, environmental, and socioeconomic conditions create barriers to the effective delivery of maternal health services. Addressing these challenges is essential to further reduce maternal and neonatal morbidity and improve reproductive health outcomes.

Ladakh's mountainous landscape, scattered settlements, and harsh climatic conditions remain major barriers to maternal healthcare. Many villages are located far from primary and secondary healthcare facilities, requiring pregnant women to travel long distances for antenatal check-ups and delivery services. During winter, heavy snowfall often blocks roads and mountain passes, delaying access to emergency obstetric care and increasing the risk of pregnancy-related complications (MoHFW, 2021). Such delays are particularly critical in obstetric

International Journal of
LAW, ARTS AND HUMANITIES
Peer Reviewed, Referred & Indexed Journal
ISSN: 3070-0523 www.zesterapublications.com Original Research Paper

emergencies where timely intervention can determine maternal and neonatal survival.A persistent shortage of obstetricians, gynaecologists, anaesthetists, paediatricians, and trained nursing staff affects the quality of maternal healthcare in Ladakh. While Primary Health Centres (PHCs) and Community Health Centres (CHCs) provide essential maternal services, complicated pregnancies often require referral to district hospitals in Leh or Kargil. Recruiting and retaining skilled healthcare professionals in remote high-altitude areas remain a significant challenge due to difficult living conditions and limited professional opportunities (National Health Mission, 2021).The high-altitude environment presents additional risks during pregnancy. Reduced oxygen availability, extreme cold, and harsh weather conditions can contribute to complications such as fetal growth restriction, low birth weight, preterm birth, and hypertensive disorders of pregnancy (Julian, 2011). These environmental factors also complicate emergency transportation and healthcare delivery, especially during adverse weather conditions.Though maternal nutrition programmes have expanded under the National Health Mission, nutritional deficiencies remain an important concern. Limited agricultural production, seasonal food shortages, and inadequate dietary diversity may contribute to iron deficiency and anaemia among women of reproductive age. Maternal anaemia increases the risk of maternal morbidity, postpartum haemorrhage, preterm delivery, and low birth weight, highlighting the need for strengthened nutritional interventions and regular screening during pregnancy (World Health Organization [WHO], 2023). Access to reproductive healthcare services, including family planning, adolescent reproductive health education, infertility counselling, and reproductive cancer screening, remains uneven across Ladakh. Women living in remote villages may experience difficulties accessing specialised reproductive healthcare because of geographical barriers and limited availability of trained healthcare providers. Strengthening community-based reproductive health services and expanding outreach programmes are essential to ensure equitable healthcare access.

Digital health technologies and telemedicine have considerable potential to improve maternal healthcare in geographically isolated regions. However, unreliable internet connectivity, inconsistent electricity supply, and limited digital infrastructure restrict the effective implementation of telemedicine services in many parts of Ladakh. Similarly, emergency referral systems, ambulance availability, and communication networks require further strengthening to improve timely management of obstetric emergencies.The challenges affecting maternal and reproductive healthcare in Ladakh are multidimensional and closely interconnected. Geographic isolation, high-altitude conditions, shortages of skilled healthcare professionals, nutritional deficiencies, and limited emergency referral systems collectively influence maternal health outcomes. Although national programmes have substantially improved service coverage, achieving equitable maternal healthcare requires context-specific strategies tailored to the environmental and healthcare realities of

Received: 08-12-2025 | Accepted Received: 02-01-2026 | Accepted: 06-02-2026 |

International Journal of
LAW, ARTS AND HUMANITIES
Peer Reviewed, Referred & Indexed Journal
ISSN: 3070-0523 www.zesterapublications.com Original Research Paper

the Himalayan region. Greater investment in healthcare infrastructure, workforce development, digital health technologies, and community-based maternal care is essential to ensure safe motherhood and universal access to reproductive healthcare.

**Table 4. Major Challenges to Maternal and Reproductive Healthcare in Ladakh**

| Challenge | Impact on Maternal Health |
|---|---|
| Geographic isolation | Delays access to antenatal care and emergency obstetric services |
| Harsh climate and seasonal road closures | Interrupts transportation and referral services |
| Shortage of skilled healthcare professionals | Limits access to specialised maternal healthcare |
| High-altitude hypoxia | Increases risk of pregnancy complications and adverse birth outcomes |
| Maternal anaemia and nutritional deficiencies | Contributes to maternal morbidity and poor neonatal outcomes |
| Limited digital health infrastructure | Restricts telemedicine and specialist consultations |
| Inadequate emergency referral systems | Delays timely management of obstetric emergencies |

*Source: Secondary Data*

**7. Government Policies and Programmes: Achievements and Gaps**

The Government of India has implemented several initiatives to improve maternal and reproductive health, including the National Health Mission (NHM), Janani Suraksha Yojana (JSY), Janani Shishu Suraksha Karyakram (JSSK), Pradhan Mantri Surakshit Matritva Abhiyan (PMSMA), Ayushman Bharat, Anaemia Mukt Bharat (AMB), and the Rashtriya Kishor Swasthya Karyakram (RKSK). These programmes aim to increase antenatal care coverage, promote institutional deliveries, improve maternal nutrition, strengthen reproductive healthcare services, and reduce maternal and neonatal mortality (MoHFW, 2021).

In Ladakh, the implementation of these programmes has contributed to improved maternal health indicators. The high rate of institutional deliveries and antenatal care coverage reported in NFHS-5 reflects increased access to skilled maternal healthcare and greater awareness among pregnant women (IIPS & MoHFW, 2021). Frontline health workers, including Accredited Social Health Activists (ASHAs), Auxiliary Nurse Midwives (ANMs), and Anganwadi Workers (AWWs), have played an important role in promoting early pregnancy registration, immunization, nutrition counselling, and institutional delivery. Despite these achievements, several implementation challenges remain. Geographical isolation, harsh weather conditions, shortages of specialist healthcare professionals, and limited emergency obstetric services reduce the effectiveness of national programmes in remote areas. In addition, access to comprehensive reproductive healthcare, telemedicine, and specialised maternal services remains uneven across the Union Territory. These challenges highlight the need for region-specific strategies that strengthen healthcare

Received: 08-12-2025 | Accepted Received: 02-01-2026 | Accepted: 06-02-2026 |

Zestera PUBLICATIONS ISSN: 3070-0523
International Journal of
LAW, ARTS AND HUMANITIES
Peer Reviewed, Referred & Indexed Journal
www.zesterapublications.com
Original Research Paper

infrastructure, improve referral systems, and expand digital health services to ensure equitable maternal healthcare in Ladakh.

**Table 5. Major Maternal Health Programmes and their Relevance to Ladakh**

| Programme | Objective | Relevance to Ladakh |
|---|---|---|
| National Health Mission (NHM) | Strengthen primary healthcare | Improves maternal and child health services |
| Janani Suraksha Yojana (JSY) | Promote institutional deliveries | Increases skilled birth attendance |
| Janani Shishu Suraksha Karyakram (JSSK) | Free maternal and newborn care | Reduces financial barriers |
| PMSMA | Quality antenatal care | Supports early detection of high-risk pregnancies |
| Ayushman Bharat | Universal health coverage | Expands access to healthcare services |
| Anaemia Mukt Bharat | Reduce anaemia | Improves maternal nutrition |
| RKSK | Adolescent health | Promotes reproductive health awareness |

*Source: Secondary Data*

## 8. Critical Synthesis of Current Evidence

The available evidence indicates that maternal and reproductive health in Ladakh has improved considerably over the past decade through increased antenatal care utilization, institutional deliveries, and implementation of national maternal health programmes. The findings from NFHS-5 (2019–21) demonstrate encouraging progress in key maternal health indicators, reflecting the positive impact of initiatives under the National Health Mission (IIPS & MoHFW, 2021). However, despite these improvements, geographical isolation, high-altitude hypoxia, workforce shortages, and limited emergency obstetric services continue to pose significant challenges.

Most published studies on maternal health in Ladakh focus on service utilization and health programme implementation, whereas evidence on high-altitude pregnancy, maternal nutrition, mental health, and long-term reproductive outcomes remains limited. Furthermore, many studies are descriptive and based on secondary data, highlighting the need for more robust epidemiological and longitudinal research. Although findings from other high-altitude regions such as Nepal and the Andes provide useful insights, they cannot be directly generalized to Ladakh due to differences in population characteristics, healthcare systems, and environmental conditions (Julian, 2011; Moore et al., 2011). Therefore, locally generated evidence is essential for developing context-specific maternal health policies and interventions.

Received: 08-12-2025 | Accepted Received: 02-01-2026 | Accepted: 06-02-2026 |

International Journal of
LAW, ARTS AND HUMANITIES
Peer Reviewed, Referred & Indexed Journal
ISSN: 3070-0523 www.zesterapublications.com Original Research Paper

**9. Research Gaps**

Although maternal and reproductive health has been studied in Himalayan regions, significant research gaps remain in Ladakh. Limited evidence is available on the effects of high-altitude conditions on maternal and neonatal health, maternal nutrition and anaemia, and maternal mental health. Research on reproductive healthcare among remote and nomadic populations is also scarce. Furthermore, few studies have examined the role of telemedicine, seasonal accessibility, and climate-related challenges in maternal healthcare delivery. Addressing these gaps through comprehensive research is essential for developing evidence-based policies and improving reproductive and maternal health services in Ladakh.

**10. Policy Recommendations**

Improving reproductive and maternal health in Ladakh requires strengthening primary healthcare services, ensuring timely access to antenatal and emergency obstetric care, and expanding outreach services in remote areas. Efforts should focus on improving maternal nutrition, increasing iron and folic acid supplementation, promoting family planning awareness, and strengthening routine screening for high-risk pregnancies and anaemia. The use of telemedicine and mobile health services can further enhance access to quality maternal healthcare in geographically isolated communities.

**11. Conclusion**

Maternal and reproductive health in Ladakh has improved considerably through increased antenatal care coverage, institutional deliveries, and implementation of national maternal health programmes. Nevertheless, the region continues to face significant challenges related to high-altitude geography, limited healthcare infrastructure, shortages of skilled healthcare professionals, maternal anaemia, and barriers to accessing emergency obstetric care. These factors underscore the need for region-specific healthcare strategies that consider the unique environmental and demographic characteristics of Ladakh. Strengthening healthcare infrastructure, improving maternal nutrition, expanding telemedicine, and developing high-altitude-specific clinical guidelines are essential to achieving equitable maternal healthcare. In addition, greater investment in locally driven research and evidence-based policymaking will contribute to improved maternal and neonatal outcomes. A coordinated approach involving government agencies, healthcare providers, researchers, and local communities will be critical for ensuring safe motherhood and advancing reproductive health in this geographically unique Himalayan region.

**References**

International Institute for Population Sciences (IIPS), & Ministry of Health and Family Welfare. (2021). *National Family Health Survey (NFHS-5), 2019–21: Union Territory Fact Sheet – Ladakh*. Government of India.

International Institute for Population Sciences (IIPS), & Ministry of Health and Family Welfare. (2026). *National Family Health Survey (NFHS-6), 2023–24: Union Territory Fact Sheet – Ladakh*. Government of India.

Julian, C. G. (2011). High altitude during pregnancy. *Clinics in Chest Medicine,*

Received: 08-12-2025 | Accepted Received: 02-01-2026 | Accepted: 06-02-2026 |

*32*(1), 21–31. https://doi.org/10.1016/j.ccm.2010.10.008

Keyes, L. E., Armaza, J. F., Niermeyer, S., Vargas, E., Young, D. A., & Moore, L. G. (2003). Intrauterine growth restriction, preeclampsia, and intrauterine mortality at high altitude in Bolivia. *Pediatric Research, 54*(1), 20–25. https://doi.org/10.1203/01.PDR.0000069846.64389.DC

Ministry of Health and Family Welfare. (2021). *National Health Mission: Framework for Implementation*. Government of India.

Moore, L. G., Charles, S. M., & Julian, C. G. (2011). Humans at high altitude: Hypoxia and fetal growth. *Respiratory Physiology & Neurobiology, 178*(1), 181–190. https://doi.org/10.1016/j.resp.2011.04.017

Office of the Registrar General of India. (2022). *Special Bulletin on Maternal Mortality in India 2018–2020*. Government of India.

Srivastava, A., Avan, B. I., Rajbangshi, P., & Bhattacharyya, S. (2015). Determinants of women's satisfaction with maternal health care: A review of literature from developing countries. *BMC Pregnancy and Childbirth, 15*, Article 97. https://doi.org/10.1186/s12884-015-0525-0

United Nations. (2015). *Transforming our world: The 2030 agenda for sustainable development*. United Nations.

World Health Organization. (2022). *WHO recommendations on antenatal care for a positive pregnancy experience*. World Health Organization.

World Health Organization. (2023). *Trends in maternal mortality 2000–2020: Estimates by WHO, UNICEF, UNFPA, World Bank Group and UNDESA/Population Division*. World Health Organization.